Table S1. Diffusing protein parameters for spatial-stochastic simulations.

Molecular species	Initial number	Diffusion rate	Estimated size (including GFP)	Reference
GFP	-	-	(W x L) 2.4 x 4.2 nm	[1]
TetR-GFP	5000 @ nucleoplasm	1.9 $\mu m^2/s$	4 nm (8 nm)	[2]
NPC (Nup49-GFP)	150 @ periplasm	0.2 $\mu m^2/s$	98 nm	[3]
Nsg1-GFP	1900 @ ONM	0.3 $\mu m^2/s$	~ 4.2 nm (8 nm)	[4]
GFP-Src1	2100 @ INM	0.3 $\mu m^2/s$	~ 6 nm (10 nm)	[4]

References

1. Ormö M, Cubitt AB, Kallio K, Gross LA, Tsien RY, et al. (1996) Crystal structure of the Aequorea victoria green fluorescent protein. Science 273: 1392-1395.
2. Hinrichs W, Kisker C, Duvel M, Muller A, Tovar K, et al. (1994) Structure of the Tet repressor-tetracycline complex and regulation of antibiotic resistance. Science 264: 418-420.
3. Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, et al. (2007) The molecular architecture of the nuclear pore complex. Nature 450: 695-701.
4. Erickson HP (2009) Size and shape of protein molecules at the nanometer level determined by sedimentation, gel filtration, and electron microscopy. Biol Proced Online 11: 32-51.